

CURVATURE FLOW TO NIRENBERG PROBLEM

MINCHUN HONG, LI MA

ABSTRACT. In this note, we study the curvature flow to Nirenberg problem on S^2 with non-negative nonlinearity. This flow was introduced by Brendle and Struwe. Our result is that the Nirenberg problems has a solution provided the prescribed non-negative Gaussian curvature f has its positive part, which possesses non-degenerate critical points such that $\Delta_{S^2} f > 0$ at the saddle points.

Mathematics Subject Classification 2000: 53Cxx, 35Jxx

Keywords: Nirenberg problem, curvature flow, non-negative nonlinearity

1. INTRODUCTION

In the interesting paper [7], M.Struwe studied a heat flow method to the Nirenberg problem on S^2 . This kind of heat flow in conformal geometry was considered by S.Brendle in [2]. Given the Riemannian metric g on S^2 with Gaussian curvature K . Using the well-known Gauss-Bonnet formula

$$\int_{S^2} K dv_g = 4\pi,$$

we know that K has to be positive some where. This gives a necessary condition for the Nirenberg problem on S^2 . Assuming the prescribed curvature function f being positive on S^2 , the heat flow for the Nirenberg problem S^2 is a family of metrics of the form $g = e^{2u(x,t)}c$ satisfying

$$(1) \quad u_t = \alpha f - K, \quad x \in S^2, \quad t > 0,$$

where c is the standard spherical metric on S^2 , $u : S^2 \times (0, T) \rightarrow R$, and $\alpha = \alpha(t)$ is defined by

$$(2) \quad \alpha \int_{S^2} f dv_g = 4\pi.$$

Here dv_g is the area element with respect to the metric g . It is easy to see that

$$\alpha_t \int_{S^2} f dv_g = 2\alpha \int_{S^2} (K - \alpha f) f dv_g.$$

M.Struwe can show that the flow exists globally, furthermore, the flow converges at infinity provided f is positive and possesses non-degenerate critical

Date: July. 12th, 2008.

* Li Ma: Corresponding author. The research is partially supported by the National Natural Science Foundation of China 10631020 and SRFDP 20060003002 .

points such that $\Delta_{S^2} f > 0$ at the saddle points. Here $\Delta_{S^2} := \Delta$ is the Analyst's Laplacian on the standard 2-sphere (S^2, c) . Recall that $\int_{S^2} dv_c = 4\pi$. The purpose of this paper is to relax his assumption by allowing the function f to have zeros.

Since we have

$$K = e^{2u}(-\Delta u + 1),$$

the equation (1) define a nonlinear parabolic equation for u , and the flow exists at least locally for any initial data $u|_{t=0} = u_0$. Clearly, we have

$$\partial_t \int_{S^2} dv_g = 2 \int_{S^2} S^2 u_t dv_g = 0.$$

We shall assume that the initial data u_0 satisfies the condition

$$(3) \quad \int f e^{2u} dv_c > 0.$$

We shall show that this property is preserved along the flow. It is easy to compute that

$$(4) \quad K_t = -2u_t K - \Delta_g u_t = 2K(K - \alpha f) + \Delta_g(K - \alpha f),$$

where $\Delta_g = e^{-2u} \Delta$. Using (4), we can compute the growth rate of the Calabi energy $\int_{S^2} |K - \alpha f|^2 dv_g$.

Our main result is following

Theorem 1. *Let f be a positive somewhere, non-negative smooth function on S^2 with only non-degenerate critical points on the its positive part f_+ . Suppose that there are at least two local positive maxima of f , and at all positive valued saddle points q of f there holds $\Delta_{S^2} f(q) > 0$. Then f is the Gaussian curvature of the conformal metric $g = e^{2u} c$ on S^2 .*

Note that this result is an extension of the famous result of Chang-Yang [3] where only positive f has been considered. A similar result for Q-curvature flow has been obtained in [9].

For simplifying notations, we shall use the conventions that $dc = \frac{dv_c}{4\pi}$ and $\bar{u} = \bar{u}(t)$ defined by

$$\int_{S^2} (u - \bar{u}) dv_c = 0.$$

2. BASIC PROPERTIES OF THE FLOW

Recall the following result of Onofri-Hong [5] that

$$(5) \quad \int_{S^2} (|\nabla u|^2 + 2u) dc \geq \log \left(\int_{S^2} e^{2u} dc \right) = 0,$$

where $|\nabla u|^2$ is the norm of the gradient of the function u with respect to the standard metric c . Here we have used the fact that $\int_{S^2} e^{2u} dc = 1$ along the flow (1).

We show that this condition is preserved along the flow (1). In fact, letting

$$E(u) = \int_{S^2} (|\nabla u|^2 + 2u) dc$$

be the Liouville energy of u and letting

$$E_f(u) = E(u) - \log\left(\int_{S^2} f e^{2u} dc\right)$$

be the energy function for the flow (1), we then compute that

$$(6) \quad \partial_t E_f(u) = -2 \int_{S^2} |\alpha f - K|^2 dv_g \leq 0.$$

One may see Lemma 2.1 in [7] for a proof. Hence

$$E_f(u(t)) \leq E_f(u_0), \quad t > 0.$$

After using the inequality (5) we have

$$\log(1/\int_{S^2} f e^{2u} dc) \leq E_f(u_0),$$

which implies that $\int_{S^2} f e^{2u} dv_c > 0$ and

$$e^{E_f(u_0)} \int_{S^2} e^{2u} dc \leq \int_{S^2} f e^{2u} dc.$$

Note also that $\int_{S^2} f e^{2u} dc = 1/\alpha(t)$. Hence,

$$\alpha(t) \leq \frac{1}{e^{E_f(u_0)}}.$$

Using the definition of $\alpha(t)$ we have

$$\alpha(t) \geq \frac{1}{\max_{S^2} f}.$$

We then conclude that $\alpha(t)$ is uniformly bounded along the flow, i.e.,

$$(7) \quad \frac{1}{\max_{S^2} f} \leq \alpha(t) \leq \frac{1}{e^{E_f(u_0)}}.$$

We shall use this inequality to replace (26) in [7] in the study of the normalized flow, which will be defined soon following the work of M.Struwe [7]. If we have a global flow, then using (6) we have

$$2 \int_0^\infty dt \int_{S^2} |\alpha f - K|^2 dv_g \leq 4\pi(E_f(u_0) + \log \max_{S^2} f).$$

Hence we have a suitable sequence $t_l \rightarrow \infty$ with associated metrics $g_l = g(t_l)$ and $\alpha(t_l) \rightarrow \alpha > 0$, and letting $K_l = K(g_l)$, such that

$$\int_{S^2} |K_l - \alpha f|^2 \rightarrow 0, \quad (t_l \rightarrow \infty).$$

Therefore, once we have a limiting metric g_∞ of the sequence of the metrics g_l , it follows that $K(g_\infty) = \alpha f$. After a re-scaling, we see that f is the

Gaussian curvature of the metric βg_∞ for some $\beta > 0$, which implies our Theorem 1.

3. NORMALIZED FLOW AND THE PROOF OF THEOREM 1

We now introduce a normalized flow. For the given flow $g(t) = e^{2u(t)}c$ on S^2 , there exists a family of conformal diffeomorphisms $\phi = \phi(t) : S^2 \rightarrow S^2$, which depends smoothly on the time variable t , such that for the metrics $h = \phi^*g$, we have

$$\int_{S^2} x dv_h = 0, \text{ for all } t \geq 0.$$

Here $x = (x^1, x^2, x^3) \in S^2 \subset R^3$ is a position vector of the standard 2-sphere. Let

$$v = u \circ \phi + \frac{1}{2} \log(\det(d\phi)).$$

Then we have $h = e^{2v}c$. Using the conformal invariance of the Liouville energy [3], we have

$$E(v) = E(u),$$

and furthermore,

$$\text{Vol}(S^2, h) = \text{Vol}(S^2, g) = 4\pi, \text{ for all } t \geq 0.$$

Assume $u(t)$ satisfies (1) and (2). Then we have the uniform energy bounds

$$0 \leq E(v) \leq E(u) = E_f(u) + \log\left(\int_{S^2} f e^{2u} dc\right) \leq E_f(u_0) + \log(\max_{S^2} f).$$

Using Jensen's inequality we have

$$2\bar{v} := \int_{S^2} 2v dc \leq \log\left(\int_{S^2} e^{2v} dc\right) = 0.$$

Using this we can obtain the uniform H^1 norm bounds of v for all $t \geq 0$ that

$$\sup_t |v(t)|_{H^1(S^2)} \leq C.$$

See the proof of Lemma 3.2 in [7]. Using the Aubin-Moser-Trudinger inequality [1] we further have

$$\sup_t \int_{S^2} e^{2pv(t)} dc \leq C(p)$$

for any $p \geq 1$.

Note that

$$v_t = u_t \circ \phi + \frac{1}{2} e^{-2v} \text{div}_{S^2}(\xi e^{2v})$$

where $\xi = (d\phi)^{-1}\phi_t$ is the vector field on S^2 generating the flow $(\phi(t))$, $t \geq 0$, as in [7], formula (17), with the uniform bound

$$|\xi|_{L^\infty(S^2)} \leq C \int_{S^2} |\alpha f - K|^2 dv_g.$$

With the help of this bound, we can show (see Lemma 3.3 in [7]) that for any $T > 0$, it holds

$$\sup_{0 \leq t < T} \int_{S^2} e^{4|u(t)|} dc < +\infty.$$

Following the method of M.Struwe [6] (see also Lemma 3.4 in [7]) and using the bound (7) and the growth rate of α , we can show that

$$\int_{S^2} |\alpha f - K|^2 dv_g \rightarrow 0$$

as $t \rightarrow \infty$. Once getting this curvature decay estimate, we can come to consider the concentration behavior of the metrics $g(t)$. Following [6], we show that

Lemma 2. *Let (u_l) be a sequence of smooth functions on S^2 with associated metrics $g_l = e^{2u_l}c$ with $\text{vol}(S^2, g_l) = 4\pi$, $l = 1, 2, \dots$. Suppose that there is a smooth function K_∞ , which is positive somewhere and non-negative in S^2 such that*

$$|K(g_l) - K_\infty|_{L^2(S^2, g_l)} \rightarrow 0$$

as $l \rightarrow \infty$. Let $h_l = \phi_l^* g_l = e^{2v_l}c$ be defined as before. Then we have either

- 1) for a subsequence $l \rightarrow \infty$ we have $u_l \rightarrow u_\infty$ in $H^2(S^2, c)$, where $g_\infty = e^{2u_\infty}c$ has Gaussian curvature K_∞ , or
- 2) there exists a subsequence, still denoted by (u_l) and a point $q \in S^2$ with $K_\infty(q) > 0$, such that the metrics g_l has a measure concentration that

$$dv_{g_l} \rightarrow 4\pi\delta_q$$

weakly in the sense of measures, while $h_l \rightarrow c$ in $H^2(S^2, c)$ and in particular, $K(h_l) \rightarrow 1$ in $L^2(S^2)$. Moreover, in the latter case the conformal diffeomorphisms ϕ_l weakly converges in $H^1(S^2)$ to the constant map $\phi_\infty = q$.

Proof. The case 1) can be proved as Lemma 3.5 in [7]. So we need only to prove the case 2). As in [7], we choose $q_l \in S^2$ and radii $r_l > 0$ such that

$$\sup_{q \in S^2} \int_{B(q, r_l)} |K(g_l)| dv_{g_l} \leq \int_{B(q_l, r_l)} |K(g_l)| dv_{g_l} = \pi,$$

where $B(q, r_l)$ is the geodesic ball in (S^2, g_l) . Then we have $r_l \rightarrow 0$ and we may assume that $q_l \rightarrow q$ as $l \rightarrow \infty$. For each l , we introduce ϕ_l as in Lemma 3.5 in [7] so that the functions

$$\hat{u}_l = u_l \circ \phi_l + \frac{1}{2} \log(\det(d\phi_l))$$

satisfy the conformal Gaussian curvature equation

$$-\Delta_{R^2} \hat{u}_l = \hat{K}_l e^{2\hat{u}_l}, \text{ on } R^2,$$

where $\hat{K}_l = K(g_l) \circ \phi$ and Δ_{R^2} is the Laplacian operator of the standard Euclidean metric g_{R^2} . Note that for $\hat{g}_l = \phi^* g_l = e^{2\hat{u}_l} g_{R^2}$, we have

$$\text{Vol}(R^2, \hat{g}_l) = \text{Vol}(S^2, g_l) = 4\pi.$$

Arguing as in [7], we can conclude a convergent subsequence $\hat{u}_l \rightarrow \hat{u}_\infty$ in $H_{loc}^2(R^2)$ where $\hat{u}_\infty$ satisfies the Liouville equation

$$-\Delta_{R^2} \hat{u}_\infty = \hat{K}_\infty(q) e^{2\hat{u}_\infty}, \text{ on } R^2,$$

with the finite volume $\int_{R^2} e^{2\hat{u}_\infty} dz \leq 4\pi$.

We remark that blow up point only occurs at point where $K_\infty > 0$ provided we allow K_∞ to change sign. We only need to exclude the case when $K_\infty(q) \leq 0$. It is clear that $K_\infty(q) < 0$ can not occur since there is no such a solution on the whole plane R^2 [8]. If $K_\infty(q) = 0$, then $\hat{u} := \hat{u}_\infty$ is a harmonic function in R^2 . Let $\bar{u}(r)$ be the average of u on the circle $\partial B_r(0) \subset R^2$. Then we have

$$\Delta_{R^2} \bar{u} = 0.$$

Hence $\bar{u} = A + B \log r$ for some constants A and B , where $r = |x|$. Since $\bar{u}$ is a continuous function on $[0, \infty)$, we have $\bar{u} = A$, which is impossible since we have by Jensen's inequality that

$$2\pi \int_0^\infty e^{2\bar{u}(r)} r dr \leq \int_{R^2} e^{2\hat{u}_\infty} dz \leq 4\pi.$$

We now have $K_\infty(q) > 0$. Recall that we have assumed $K_\infty \geq 0$. So we can follow the proof of Lemma 3.5 in [7]. In fact, using the classification result of Chen-Li [4] we know that $\hat{u}_\infty$ can be obtained from stereographic projection $S^2 \rightarrow R^2$ with

$$\int_{R^2} K_\infty(q) e^{2\hat{u}_\infty} dz = 4\pi.$$

Thus for any large $R > 0$, we have error $\circ(1) \rightarrow 0$ as $l \rightarrow \infty$ such that

$$4\pi = \int_{R^2} K_\infty(q) e^{2\hat{u}_\infty} dz \leq \int_{B_R(q)} K_l dv_l + \circ(1) \leq \int_{B_R(q)} |K_l| dv_l + \circ(1).$$

Then the remaining part can be derived as in [7]. We refer to [7] for the full proof. □

With this understanding, we can do the same finite-dimensional dynamics analysis as in section 4 in [7]. Then arguing as in section 5 in [7] we can prove Theorem 1. By now the argument is well-known, we omit the detail and refer to [7] for full discussion.

REFERENCES

- [1] Aubin, Thierry: *Some nonlinear problems in Riemannian geometry* (Springer Monographs in Mathematics). Springer, Berlin 1998 MR1636569 (99i:58001)
- [2] S.Brendle, *Global existence and convergence for a higher order flow in conformal geometry*, Ann. math., 158(2003)323-343.
- [3] S.Y.A.Chang, P.Yang, *Prescribing Gaussian curvature on S^2* , Acta Mathematica, 159(1987)215-296.
- [4] W.Chen, C.Li, *Classification of solutions of some nonlinear equations*, Duke Math. Journal, 63(1991)615-623.

- [5] E.Onofri, *On the positivity of the effective action in a theory of random surfaces*, Comm. Math. Phys., 86(1982)321-326.
- [6] M.Struwe, *Curvature flows on surfaces*, Annali Sc. Norm. Sup. Pisa, Ser. V, 1(2002)247-274.
- [7] M.Struwe, *A flow approach to Nirenberg's problem*, Duke Math. J. 128(2005)19-64.
- [8] L.Ma, *Three remarks on mean field equations*, preprint, 2008.
- [9] L.Ma, *Q-curvature flow with indefinite nonlinearity*, preprint, 2008.

LI MA, DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING 100084, CHINA

E-mail address: `lma@math.tsinghua.edu.cn`

M.C.HONG, DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF QUEENSLAND, BRISBANE, QUEENSLAND, AUSTRALIA

E-mail address: `hong@maths.uq.edu.au`